arXiv:1606.02161v1 [math.DS] 7 Jun 2016

The numbers of distinct and repeated squares and cubes in the Tribonacci sequence

Huang Yuke[1] Wen Zhiying[2]

Abstract: The Tribonacci sequence $\mathbb{T}$ is the fixed point of the substitution $\sigma(a,b,c)=(ab,ac,a)$. The main result is twofold: (1) we give the explicit expressions of the numbers of distinct squares and cubes in $\mathbb{T}[1,n]$ (the prefix of $\mathbb{T}$ of length n); (2) we give algorithms for counting the number of repeated squares and cubes in $\mathbb{T}[1,n]$ for all n; then get explicit expressions for some special n such as $n=t_m$ (the Tribonacci number).

Key words the Tribonacci sequence; square; cube; algorithm; gap sequence.

2010 MR Subject Classification 11B85; 68Q45

1 Introduction

Let $\mathcal{A}=\{a,b,c\}$ be a three-letter alphabet. The Tribonacci sequence $\mathbb{T}$ is the fixed point of the substitution $\sigma(a,b,c)=(ab,ac,a)$. As a natural generalization of the Fibonacci sequence, the Tribonacci sequence has been studied extensively by many authors, see [3, 6, 7].

Let ω be a factor of $\mathbb{T}$, denoted by $\omega\prec\mathbb{T}$. Let ω_p be the p-th occurrence of ω. If the factor ω and integer p such that $\omega_p\omega_{p+1}\prec\mathbb{T}$, we call $\omega_p\omega_{p+1}$ a square of $\mathbb{T}$. We call $\omega\omega$ a square of $\mathbb{T}$ if there exist p such that $\omega_p\omega_{p+1}\prec\mathbb{T}$. Similarly, we call $\omega_p\omega_{p+1}\omega_{p+2}$ (resp. $\omega\omega\omega$) a cube of $\mathbb{T}$. As we know, $\mathbb{T}$ contains no fourth powers. The properties of squares and cubes are objects of a great interest in many aspects of mathematics and computer science etc.

We denote by $|\omega|$ the length of ω, and by $|\omega|_\alpha$ the number of letter α in ω, $\alpha\in\mathcal{A}$. Denote $P(\omega,p)$ the position of the last letter of ω_p. Let $\tau=x_1\cdots x_n$. For $i\le j\le n$, define $\tau[i,j]=x_ix_{i+1}\cdots x_{j-1}x_j$. By convention, denote $\tau[i]=\tau[i,i]=x_i$, $\tau[i,i-1]=\varepsilon$ (empty word). Denote $T_m=\sigma^m(a)$ for $m\ge0$, $T_{-2}=\varepsilon$, $T_{-1}=c$, then $T_0=a$, $T_1=ab$ and $T_m=T_{m-1}T_{m-2}T_{m-3}$ for $m\ge2$. Denote $t_m=|T_m|$ for $m\ge-2$, called the m-th Tribonacci number. Denote by δ_m the last letter of T_m, then $\delta_m=a$ (resp. b, c) for $m\equiv0$, (resp. 1, 2) mod 3 and $m\ge-1$.

Let $\mathbb{T}[1,n]$ be the prefix of $\mathbb{T}$ of length n. In this paper, we consider the four functions below:

$A(n)=\sharp\{\omega:\omega\omega\prec\mathbb{T}[1,n]\}$, the number of distinct squares in $\mathbb{T}[1,n]$;

$B(n)=\sharp\{(\omega,p):\omega_p\omega_{p+1}\prec\mathbb{T}[1,n]\}$, the number of repeated squares in $\mathbb{T}[1,n]$;

$C(n)=\sharp\{\omega:\omega\omega\omega\prec\mathbb{T}[1,n]\}$, the number of distinct cubes in $\mathbb{T}[1,n]$;

$D(n)=\sharp\{(\omega,p):\omega_p\omega_{p+1}\omega_{p+2}\prec\mathbb{T}[1,n]\}$, the number of repeated cubes in $\mathbb{T}[1,n]$.

In 2006, A.Glen[2] gave expressions of $A(t_m)$. In 2014, H.Mousavi and J.Shallit[6] gave expressions of $B(t_m)$ and $D(t_m)$, which they proved by mechanical way. All of these results above only consider the squares or cubes in the prefixes of some special lengths: the Tribonacci numbers. In this paper, for all $n,m\ge1$, we give: (1) explicit expressions of $A(n)$, $C(n)$, $B(t_m)$ and $D(t_m)$; (2) fast algorithms for counting $B(n)$ and $D(n)$. We also considered this problem in the Fibonacci sequence, see [4].

We denote the gap between ω_p and ω_{p+1} by $G_p(\omega)$. The sequence $\{G_p(\omega)\}_{p\ge1}$ is called the gap sequence of factor ω. Taking "square" for example, the **main difficulty** is twofold: (1) The positions of all squares are not easy to be determined. We overcome this difficulty by using the "gap sequence" property of $\mathbb{T}$, which we introduced and studied in [3]. (2) By the gap sequence property of $\mathbb{T}$, we can find out all distinct squares in $\mathbb{T}[1,n]$. We can also count the number of occurrences of each square. So the summation of these numbers are the numbers of repeated squares in $\mathbb{T}[1,n]$. But this method

[1]School of Mathematics and Systems Science, Beihang University (BUAA), Beijing, 100191, P. R. China. E-mail address: huangyuke@buaa.edu.cn, hyg03ster@163.com (Corresponding author).

[2]Department of Mathematical Sciences, Tsinghua University, Beijing, 100084, P. R. China. E-mail address: wenzy@tsinghua.edu.cn.

is complicated. We overcome this difficulty by studying the relations among positions of each ω_p, and establishing a recursive structure, called square trees. First we list the main results as below.

Theorem A (The numbers of distinct squares, $A(n)$)**.** *$A(n) = 0$ for $n \le 7$; $A(n) = 1$ for $n = 8, 9$; $A(n) = 2$ for $10 \le n \le 13$. For $n \ge 14$, let m such that $\alpha_m \le n < \alpha_{m+1}$, then $m \ge 4$,*

$$A(n) = \begin{cases} n - \frac{1}{2}(t_m + t_{m-3} + m + 3), & \alpha_m \le n < \beta_m; \\ \frac{1}{2}(t_{m-1} + t_{m-2} + 4t_{m-3} - m - 5), & \beta_m \le n < \gamma_m; \\ n - \frac{1}{2}(t_{m-1} + 3t_{m-2} + m + 3), & \gamma_m \le n < \theta_m; \\ \frac{1}{2}(2t_{m-1} + t_{m-2} + 3t_{m-3} - m - 6), & \theta_m \le n < \alpha_{m+1}. \end{cases}$$

Here $\alpha_m = 2t_{m-1}$, $\beta_m = t_m + 2t_{m-3} - 1$, $\gamma_m = 2t_m - t_{m-1}$ and $\theta_m = \frac{3t_m + t_{m-2} - 3}{2}$.

Example. Consider $n = 65 \in [\alpha_6, \cdots, \alpha_7 - 1] = [48, \cdots, 87]$, $m = 6$. Moreover $\gamma_6 = 2t_6 - t_5 = 64 \le n < \theta_6 = \frac{3t_6 + t_4 - 3}{2} = 71$, $A(65) = 65 - \frac{1}{2}(t_5 + 3t_4 + 6 + 3) = 29$. In fact, the positions of the last letters of the 29 squares are $\{8, 10, 14, 15, 16, 19, 20, \underbrace{26, \cdots, 31}_{6}, \underbrace{35, \cdots, 38}_{4}, \underbrace{48, \cdots, 57}_{10}, 64, 65\}$. Here the number under "$\smile$" means the number of elements.

Theorem B (The numbers of repeated squares, $B(t_m)$)**.** *For $m \ge 3$,*

$$B(t_m) = \tfrac{m}{22}(9t_m - t_{m-1} - 5t_{m-2}) + \tfrac{1}{44}(-81t_m + 26t_{m-1} + 13t_{m-2}) + m + \tfrac{1}{4}.$$

Example. $B(t_5) = B(24) = \frac{5}{22}(9t_5 - t_4 - 5t_3) + \frac{1}{44}(-81t_5 + 26t_4 + 13t_3) + 5 + \frac{1}{4} = 9$.

Theorem C (The numbers of distinct cubes, $C(n)$)**.** *$C(n) = 0$ for $n \le 57$. For $n \ge 58$, let m such that $t_{m-1} + 2t_{m-4} \le n < t_m + 2t_{m-3}$, then $m \ge 7$,*

$$C(n) = \begin{cases} n - \frac{1}{2}(4t_{m-1} - t_{m-2} - 3t_{m-3} + m - 6), & n \le \frac{3t_{m-1} - t_{m-3} - 3}{2}; \\ \frac{1}{2}(t_{m-5} + t_{m-6} - m + 3), & otherwise. \end{cases}$$

Example. Consider $n = 365 \in [t_9 + 2t_6, \cdots, t_{10} + 2t_7] = [362, \cdots, 666]$, $m = 10$. Moreover $n \le \frac{3t_9 - t_7 - 3}{2} = 369$, $C(365) = 365 - \frac{1}{2}(4t_9 - t_8 - 3t_7 + 10 - 6) = 11$. In fact, the positions of the last letters of the 11 cubes are $\{58, 107, 108, 197, 198, 199, 200, 362, 363, 364, 365\}$.

Theorem D (The numbers of repeated cubes, $D(t_m)$)**.** *For $m \ge 3$,*

$$\begin{aligned} D(t_m) = &\tfrac{m}{22}(-6t_m + 8t_{m-1} + 7t_{m-2}) + \tfrac{1}{44}(-23t_m + 34t_{m-1} - 5t_{m-2}) + \tfrac{m}{6} \\ &- \tfrac{1}{4}[m \equiv 0 (\mathrm{mod}\ 3)] + \tfrac{1}{12}[m \equiv 1 (\mathrm{mod}\ 3)] + \tfrac{5}{12}[m \equiv 2 (\mathrm{mod}\ 3)]. \end{aligned}$$

Here $[P]$ is Iverson notation, and equals 1 if P holds and 0 otherwise.

Example. $D(t_8) = D(149) = \frac{8}{22}(-6t_8 + 8t_7 + 7t_6) + \frac{1}{44}(-23t_8 + 34t_7 - 5t_6) + \frac{8}{6} + \frac{5}{12} = 4$.

H.Mousavi and J.Shallit gave **Theorem B.** and **Theorem D.** in [6]. We prove them as corollaries of **Algorithm B.** and **Algorithm D.**, which are fast algorithms for counting $B(n)$ and $D(n)$ for all n, see Section 4 and 6, respectively.

2 Preliminaries of the Tribonacci sequence

We define the kernel numbers that $k_0 = 0$, $k_1 = k_2 = 1$, $k_m = k_{m-1} + k_{m-2} + k_{m-3} - 1$ for $m \ge 3$. Then $k_m = k_{m-3} + t_{m-4} = \frac{t_{m-3} + t_{m-5} + 1}{2}$ for $m \ge 3$. The kernel word with order m is defined as $K_1 = a$, $K_2 = b$, $K_3 = c$, $K_m = \delta_{m-1} T_{m-3}[1, k_m - 1]$ for $m \ge 4$. Let $Ker(\omega)$ be the maximal kernel word occurring in factor ω, then by Theorem 4.3 in [3], $Ker(\omega)$ occurs in ω only once. Moreover,

Property 2.1 (Theorem 4.11 in [3]). *$Ker(\omega_p) = Ker(\omega)_p$ for all $\omega \in \mathbb{T}$ and $p \geq 1$.*

Property 2.2 (Theorem 3.3 in [3]). *The gap sequence $\{G_p(\omega)\}_{p\geq 1}$ is the Tribonacci sequence over the alphabet $\{G_1(\omega), G_2(\omega), G_4(\omega)\}$ for all $\omega \in \mathbb{T}$.*

Property 2.3 (Property 6.1 in [5]). *For $m, p \geq 1$,*
$P(K_m, p) = pt_{m-1} + |\mathbb{T}[1, p-1]|_a(t_{m-2} + t_{m-3}) + |\mathbb{T}[1, p-1]|_b t_{m-2} + k_m - 1$.

Example. $P(K_m, 1) = k_{m+3} - 1 = \frac{t_m + t_{m-2} - 1}{2}$ for $m \geq 1$. $P(a, p) = p + |\mathbb{T}[1, p-1]|_a + |\mathbb{T}[1, p-1]|_b$, $P(b, p) = 2p + 2|\mathbb{T}[1, p-1]|_a + |\mathbb{T}[1, p-1]|_b$, $P(c, p) = 4p + 3|\mathbb{T}[1, p-1]|_a + 2|\mathbb{T}[1, p-1]|_b$ for $p \geq 1$.

Property 2.4 (Lemma 6.4. in [5]). *For $p \geq 1$, (1) $|\mathbb{T}[1, P(\alpha, p)]|_\alpha = p$ for $\alpha \in \mathcal{A}$;*
(2) $|\mathbb{T}[1, P(b, p)]|_a = P(a, p)$, $|\mathbb{T}[1, P(a, p)|_b = |\mathbb{T}[1, p-1]|_a$;
(3) $|\mathbb{T}[1, P(c, p)]|_a = P(b, p)$, $|\mathbb{T}[1, P(c, p)]|_b = P(a, p)$.

3 The number of distinct squares, $A(n)$

By Lemma 4.7, Definition 4.12 and Corollary 4.13 in [3], any factor ω with kernel K_m can be expressed uniquely as $\omega = T_{m-1}[i, t_{m-1} - 1]K_m T_m[k_m, k_m + j - 1]$, where $1 \leq i \leq t_{m-1}$ and $0 \leq j \leq t_{m-1} - 1$.

By Theorem 3.3, Corollary 3.12 and Proposition 6.7(1) in [3], $\omega_p\omega_{p+1} \prec \mathbb{T}$ has three cases.

Case 1. $G_p(K_m) = G_1(K_m)$. Here $|G_1(K_m)| = t_m - k_m$.

Since $|G_p(K_m)| = |T_{m-1}[i, t_{m-1}-1]| + |T_m[k_m, k_m+j-1]| = t_{m-1} - i + j$, $j = t_{m-2} + t_{m-3} - k_m + i$. So $0 \leq j \leq t_{m-1} - 1$ gives a range of i. Comparing this range with $1 \leq i \leq t_{m-1}$, we have $1 \leq i \leq k_{m+1} - 1$ and $m \geq 3$. Furthermore,

$$\begin{aligned}\omega =& T_{m-1}[i, t_{m-1}]T_m[1, t_{m-2} + t_{m-3} + i - 1] = T_{m-1}[i, t_{m-1}]T_{m-2}T_{m-3}T_{m-2}[1, i-1];\\ \omega\omega =& T_m[i, t_m - 1]\underline{\delta_m T_{m-1}[1, k_{m+1} - 1]}T_{m+1}[k_{m+1}, t_m + i - 1].\end{aligned}$$

Thus $K_{m+1} = \delta_m T_{m-1}[1, k_{m+1} - 1] \prec \omega\omega$. Similarly, $K_{m+2}, K_{m+3}, K_{m+4} \not\prec \omega\omega$ and $|\omega\omega| < K_{m+5}$, so K_{m+1} is the largest kernel word in $\omega\omega$, i.e. $Ker(\omega\omega) = K_{m+1}$ for $m \geq 3$. Moreover, since $|\omega| = |G_p(K_m)| + k_m$, $|\omega| = t_m$.

By analogous arguments, we have

Case 2. $G_p(K_m) = G_2(K_m)$. Here $|G_2(K_m)| = t_{m-2} + t_{m-1} - k_m$.

$$\omega\omega = T_m[i, t_{m-1} + t_{m-2} - 1]\underline{K_{m+2}}T_{m+1}[k_{m+2}, t_{m-1} + t_{m-2} + i - 1],$$

where $1 \leq i \leq k_{m+2} - 1$, $m \geq 2$, $Ker(\omega\omega) = K_{m+2}$ and $|\omega| = t_{m-2} + t_{m-1}$.

Case 3. $G_p(K_m) = G_4(K_m)$. Here $|G_4(K_m)| = t_{m-1} - k_m$.

$$\omega\omega = T_{m-1}[i, t_{m-1} - 1]\underline{K_{m+3}}T_{m+1}[k_{m+3}, t_{m-1} + i - 1],$$

where $k_m \leq i \leq t_{m-1}$, $m \geq 1$, $Ker(\omega\omega) = K_{m+3}$ and $|\omega| = t_{m-1}$.

Remark. By the three cases of squares, we have: (1) all squares in $\mathbb{T}$ are of length $2t_m$ or $2t_m + 2t_{m-1}$ for some $m \geq 0$; (2) for all $m \geq 0$, there exists a square of length $2t_m$ and $2t_m + 2t_{m-1}$ in $\mathbb{T}$. These are known results of H.Mousavi and J.Shallit, see Theorem 5 in [6].

We define three sets for $m \geq 4$,

$$\begin{cases}\langle 1, K_m, p\rangle &= \{P(\omega\omega, p) : Ker(\omega\omega) = K_m, |\omega| = t_{m-1}, \omega\omega \prec \mathbb{T}\}\\ &= \{P(K_m, p) + \frac{-t_m + 4t_{m-1} - t_{m-2} + 1}{2}, \cdots, P(K_m, p) + t_{m-1} - 1\};\\ \langle 2, K_m, p\rangle &= \{P(\omega\omega, p) : Ker(\omega\omega) = K_m, |\omega| = t_{m-4} + t_{m-3}, \omega\omega \prec \mathbb{T}\}\\ &= \{P(K_m, p) + \frac{-t_m + 4t_{m-1} - 3t_{m-2} + 1}{2}, \cdots, P(K_m, p) + t_{m-1} - t_{m-2} - 1\};\\ \langle 3, K_m, p\rangle &= \{P(\omega\omega, p) : Ker(\omega\omega) = K_m, |\omega| = t_{m-4}, \omega\omega \prec \mathbb{T}\}\\ &= \{P(K_m, p), \cdots, P(K_m, p) + \frac{-5t_m + 10t_{m-1} - t_{m-2} - 1}{2}\}.\end{cases}$$

Obviously, these sets correspond the positions $P(\omega\omega, p)$ for the three cases of squares, respectively. Each set contains some consecutive integers. Moreover $\sharp\langle 1, K_m, p\rangle = \sharp\langle 2, K_m, p\rangle = \frac{t_m - 2t_{m-1} + t_{m-2} - 1}{2}$, $\sharp\langle 3, K_m, p\rangle = \frac{-5t_m + 10t_{m-1} - t_{m-2} + 1}{2}$.

Since $P(K_m, 1) = \frac{t_m + t_{m-2} - 1}{2}$, we have $\langle 1, K_m, 1\rangle = \{2t_{m-1}, \cdots, \frac{t_m + 2t_{m-1} + t_{m-2} - 3}{2}\}$, $\langle 2, K_m, 1\rangle = \{2t_{m-1} - t_{m-2}, \cdots, \frac{t_m + 2t_{m-1} - t_{m-2} - 3}{2}\}$ and $\langle 3, K_m, 1\rangle = \{\frac{t_m + t_{m-2} - 1}{2}, \cdots, -2t_m + 5t_{m-1} - 1\}$. Obviously, $\langle j, K_m, 1\rangle$ are pairwise disjoint. Moreover, since $\max\langle 1, K_m, 1\rangle + 1 = \min\langle 3, K_{m+1}, 1\rangle$, sets $\langle 1, K_m, 1\rangle$ and $\langle 3, K_{m+1}, 1\rangle$ are consecutive. We have $\langle 1, K_m, 1\rangle \cup \langle 3, K_{m+1}, 1\rangle = \{2t_{m-1}, \cdots, t_m + 2t_{m-3} - 1\}$.

Therefore we get a chain

$$\langle 3, K_4, 1\rangle, \langle 2, K_4, 1\rangle, \langle 1, K_4, 1\rangle \cup \langle 3, K_5, 1\rangle, \cdots, \langle 2, K_m, 1\rangle, \langle 1, K_m, 1\rangle \cup \langle 3, K_{m+1}, 1\rangle, \cdots$$

Denote $a(n) = \sharp\{\omega : \omega\omega \triangleright \mathbb{T}[1, n], \omega\omega \not\prec \mathbb{T}[1, n-1]\}$, then $A(n) = \sum_{i=1}^{n} a(i)$.

Property 3.1. *For $n < 14$, $a(n) = 1$ if and only if $n \in \{8, 10\}$; for $n \geq 14$, let m such that $2t_{m-1} \leq n < 2t_m$, then $m \geq 4$ and $a(n) = 1$ if and only if*

$$n \in \{2t_{m-1}, \cdots, t_m + 2t_{m-3} - 1\} \cup \{2t_m - t_{m-1}, \cdots, \tfrac{3t_m + t_{m-2} - 3}{2}\}.$$

For $m \geq 3$, we denote $\Delta_m = \sum_{i=4}^{m} \sharp\langle j, K_i, 1\rangle$ for $j = 1, 2$, $\Theta_m = \sum_{i=4}^{m} \sharp\langle 3, K_i, 1\rangle$. So $\Delta_3 = \Theta_3 = 0$. And for $m \geq 4$, since $\sum_{i=0}^{m} t_i = \frac{t_m + t_{m+2} - 3}{2}$ and $\sum_{i=1}^{m} k_i = \frac{t_{m-2} + t_{m-3} + m}{2}$, we have $\Delta_m = \frac{t_{m-2} + t_{m-3} - m}{2}$ and $\Theta_m = \frac{t_{m-2} - t_{m-3} + 2t_{m-4} + m - 6}{2}$. For $m \geq 4$, we denote

$$\begin{cases} \alpha_m = \min\langle 1, K_m, 1\rangle = 2t_{m-1}, \ \beta_m = \max\langle 3, K_{m+1}, 1\rangle = t_m + 2t_{m-3} - 1, \\ \gamma_m = \min\langle 2, K_{m+1}, 1\rangle = 2t_m - t_{m-1}, \theta_m = \max\langle 2, K_{m+1}, 1\rangle = \frac{3t_m + t_{m-2} - 3}{2}. \end{cases}$$

By Property 3.1 and the definition of Δ_m, Θ_m,

$$\begin{cases} A(\alpha_m) = \Delta_{m-1} + \Delta_m + \Theta_m + 1 = \frac{2t_{m-2} + t_{m-3} + 3t_{m-4} - m - 3}{2}, \\ A(\beta_m) = \Delta_m + \Delta_m + \Theta_{m+1} = \frac{t_{m-1} + t_{m-2} + 4t_{m-3} - m - 5}{2}, \\ A(\gamma_m) = \Delta_m + \Delta_m + \Theta_{m+1} + 1 = A(\beta_m) + 1, \\ A(\theta_m) = \Delta_m + \Delta_{m+1} + \Theta_{m+1} = A(\alpha_{m+1}) - 1. \end{cases}$$

When $\alpha_m \leq n < \beta_m$, $A(n) = A(\alpha_m) + n - \alpha_m$; when $\beta_m \leq n < \gamma_m$, $A(n) = A(\beta_m)$; when $\gamma_m \leq n < \theta_m$, $A(n) = A(\gamma_m) + n - \gamma_m$; when $\theta_m \leq n < \alpha_{m+1}$, $A(n) = A(\theta_m)$. So we get **Theorem A**.

For $m \geq 5$, since $\theta_{m-1} = \frac{3t_{m-1} + t_{m-3} - 1}{2} \leq t_m < \alpha_m$, $A(t_m) = A(\theta_{m-1})$ by Theorem A. It is easy to check the expression holds also for $m = 3, 4$. Thus for $m = 0, 1, 2$, $A(t_m) = 0$, and

Theorem 3.2. *$A(t_m) = \frac{1}{2}(2t_{m-2} + t_{m-3} + 3t_{m-4} - m - 5)$ for $m \geq 3$.*

Remark. A.Glen gave $A(t_m)$ in Theorem 6.30 in [2], that $A(t_m) = \sum_{i=0}^{m-2}(d_i + 1) + d_{m-4} + d_{m-5} + 1$ for $m \geq 3$, where $d_{-2} = d_{-1} = -1$, $d_0 = 0$ and $d_m = \frac{t_{m+1} + t_{m-1} - 3}{2}$ for $m \geq 1$. Since $\sum_{i=0}^{m} t_i = \frac{t_m + t_{m+2} - 3}{2}$, the two expressions of $A(t_m)$ are same.

4 The number of repeated squares, $B(n)$

For $m \geq 4$ and $p \geq 1$, we consider the vectors

$$\begin{cases} \Gamma_{1,m,p} = [P(K_m, p) + t_{m-1} - t_{m-2}, \cdots, P(K_m, p) + t_{m-1} - 1]; \\ \Gamma_{2,m,p} = [P(K_m, p) - t_m + 2t_{m-1}, \cdots, P(K_m, p) + t_{m-1} - t_{m-2} - 1]; \\ \Gamma_{3,m,p} = [P(K_m, p), \cdots, P(K_m, p) - t_m + 2t_{m-1} - 1]. \end{cases}$$

Obviously, $\max\Gamma_{2,m,p}+1=\min\Gamma_{1,m,p}$ and $\max\Gamma_{3,m,p}+1=\min\Gamma_{2,m,p}$. Moreover $|\Gamma_{1,m,p}|=t_{m-2}$, $|\Gamma_{2,m,p}|=t_{m-3}$ and $|\Gamma_{3,m,p}|=t_{m-4}$. Using Property 2.4, comparing minimal and maximal elements in these sets below, we have

$$\begin{cases}\Gamma_{1,m,p}=[\Gamma_{3,m-1,P(a,p)+1},\Gamma_{2,m-1,P(a,p)+1},\Gamma_{1,m-1,P(a,p)+1}], & m\geq 5;\\ \Gamma_{2,m,p}=[\Gamma_{3,m-2,P(b,p)+1},\Gamma_{2,m-2,P(b,p)+1},\Gamma_{1,m-2,P(b,p)+1}], & m\geq 6;\\ \Gamma_{3,m,p}=[\Gamma_{3,m-3,P(c,p)+1},\Gamma_{2,m-3,P(c,p)+1},\Gamma_{1,m-3,P(c,p)+1}], & m\geq 7.\end{cases}$$

Thus we establish recursive relations for any $\Gamma_{1,m,p}$ ($m\geq 5$), $\Gamma_{2,m,p}$ ($m\geq 6$) and $\Gamma_{3,m,p}$ ($m\geq 7$).

It is easy to check that $\langle 1,K_m,p\rangle$ (resp. $\langle 2,K_m,p\rangle$, $\langle 3,K_m,p\rangle$) contains the several maximal (resp. maximal, minimal) elements of $\Gamma_{1,m,p}$ (resp. $\Gamma_{2,m,p}$, $\Gamma_{3,m,p}$). Thus we get recursive relations of the positions of repeated squares in $\mathbb{T}$, called square trees. Here we denote $P(\alpha,p)+1$ by $\hat{\alpha}$ for short, $\alpha\in\{a,b,c\}$. For $i\in\{1,2,3\}$, π_i is a substitution over $\{\langle j,K_m,p\rangle : m\geq 4+j,p\geq 1\}$ that

$$\begin{cases}\pi_1\langle 1,K_m,p\rangle=\langle 3,K_{m-1},\hat{a}\rangle\cup\langle 2,K_{m-1},\hat{a}\rangle\cup\langle 1,K_{m-1},\hat{a}\rangle, & m\geq 5;\\ \pi_2\langle 2,K_m,p\rangle=\langle 3,K_{m-2},\hat{b}\rangle\cup\langle 2,K_{m-2},\hat{b}\rangle\cup\langle 1,K_{m-2},\hat{b}\rangle, & m\geq 6;\\ \pi_3\langle 3,K_m,p\rangle=\langle 3,K_{m-3},\hat{c}\rangle\cup\langle 2,K_{m-3},\hat{c}\rangle\cup\langle 1,K_{m-3},\hat{c}\rangle, & m\geq 7.\end{cases}$$

On the other hand, for $m\geq 4$ and $j\in\{1,2,3\}$, each $\langle j,K_m,1\rangle$ belongs to the square trees. Moreover $\langle j,K_m,\hat{a}\rangle$ (resp. $\langle j,K_m,\hat{b}\rangle$, $\langle j,K_m,\hat{c}\rangle$) is subset of $\pi_1\langle 1,K_{m+1},p\rangle$ (resp. $\pi_2\langle 2,K_{m+2},p\rangle$, $\pi_3\langle 3,K_{m+3},p\rangle$). Notice that $\mathbb{N}=\{1\}\cup\{P(a,p)+1\}\cup\{P(b,p)+1\}\cup\{P(c,p)+1\}$, the square trees contain all $\langle j,K_m,p\rangle$, i.e. the positions of all squares in $\mathbb{T}$. Figure 1 shows some examples.

By the square trees, we have the relation between the number of squares ending at position $\Gamma_{j,m,p}[i]$ and $\Gamma_{j,m,1}[i]$, see Property 4.1. Figure 2 shows the relation: in the tree with root $\langle 1,K_6,1\rangle$, the branch from node $\langle 1,K_5,2\rangle$ is the graph embedding of the tree with root $\langle 1,K_5,1\rangle$.

Property 4.1. *For $j=1,2,3$, $m\geq 4$ and $p\geq 1$,*

$$\{\omega:\omega\omega\triangleright\mathbb{T}[1,\Gamma_{j,m,1}[i]]\}=\{\omega:\omega\omega\triangleright\mathbb{T}[1,\Gamma_{j,m,p}[i]],Ker(\omega)=K_h,4\leq h\leq m\}.$$

Here $1\leq i\leq k_m-1$ for $j=1,2$; $1\leq i\leq t_{m-4}-k_{m-3}+1$ for $j=3$.

Example. Let $j=1$, $m=5$, $p=3$, $i=2$. All squares ending at position $\Gamma_{1,5,1}[2]=27$ are $\{\omega\omega,\varpi\varpi\}$, where $\omega=abacaba$ and $\varpi=bacabaabacaba$. All squares ending at position $\Gamma_{1,5,3}[2]=71$ are $\{\omega\omega,\varpi\varpi,\mu\mu\}$, where $\mu=abacababacabaabacaba$. Since $Ker(\omega\omega)=aa=K_4$, $Ker(\varpi\varpi)=bab=K_5$ and $Ker(\mu\mu)=aabacabaa=K_7$, only $\{\omega\omega,\varpi\varpi\}$ are squares with kernel K_h, $4\leq h\leq 5$.

Taking $p=1$, we have

$$\begin{cases}\Gamma_{1,m,1}=[\frac{t_m+2t_{m-1}-t_{m-2}-1}{2},\cdots,\frac{t_m+2t_{m-1}+t_{m-2}-3}{2}];\\ \Gamma_{2,m,1}=[\frac{-t_m+4t_{m-1}+t_{m-2}-1}{2},\cdots,\frac{t_m+2t_{m-1}-t_{m-2}-3}{2}];\\ \Gamma_{3,m,1}=[\frac{t_m+t_{m-2}-1}{2},\cdots,\frac{-t_m+4t_{m-1}+t_{m-2}-3}{2}].\end{cases}$$

Denote $b(n)=\sharp\{(\omega,p):\omega_p\omega_{p+1}\triangleright\mathbb{T}[1,n]\}$ the number of squares ending at position n. By Property 4.2 below, we can calculate $b(n)$, and obversely calculate $B(n)$ by $B(n)=\sum_{i=1}^{n}b(i)$.

Property 4.2. *$b(i)=0$ for $i\leq 7$, $b([8])=[1]$, $b([9,10])=[0,1]$, $b([11,\cdots,14])=[0,0,0,1]$, $b([15,16])=[1,1]$, $b([17,\cdots,20])=[0,0,1,1]$, $b([28,\cdots,31])=[1,1,1,1]$,*

$$\begin{cases}b(\Gamma_{1,m,1})=[b(\Gamma_{3,m-1,1}),b(\Gamma_{2,m-1,1}),b(\Gamma_{1,m-1,1})]+[\underbrace{0,\cdots,0}_{t_{m-2}-k_m+1},\underbrace{1,\cdots,1}_{k_m-1}] \text{ for } m\geq 5;\\ b(\Gamma_{2,m,1})=[b(\Gamma_{3,m-2,1}),b(\Gamma_{2,m-2,1}),b(\Gamma_{1,m-2,1})]+[\underbrace{0,\cdots,0}_{t_{m-3}-k_m+1},\underbrace{1,\cdots,1}_{k_m-1}] \text{ for } m\geq 6;\\ b(\Gamma_{3,m,1})=[b(\Gamma_{3,m-3,1}),b(\Gamma_{2,m-3,1}),b(\Gamma_{1,m-3,1})]+[\underbrace{1,\cdots,1}_{t_{m-4}-k_{m-3}+1},\underbrace{0,\cdots,0}_{k_{m-3}-1}] \text{ for } m\geq 7.\end{cases}$$



Figure 1: (a)-(c) are square trees from roots $\langle 1, K_6, 1\rangle$, $\langle 2, K_7, 1\rangle$, $\langle 3, K_8, 1\rangle$, respectively.



Figure 2: An example of the graph embedding in square trees.

Obversely we can calculate $B(n)$ by $B(n)=\sum_{i=1}^{n}b(i)$. But when n large, the method is complicated. Now we turn to give a fast algorithm. Denote the sum of all elements in vector $b(\Gamma_{j,m,1})$ by $\sum b(\Gamma_{j,m,1})$ for $j\in\{1,2,3\}$. Denote $\Phi_m=\sum b(\Gamma_{3,m,1})+\sum b(\Gamma_{2,m,1})+\sum b(\Gamma_{1,m,1})$. The immediately corollaries of Property 4.2 are $\sum b(\Gamma_{1,m,1})=\Phi_{m-1}+k_m-1$, $\sum b(\Gamma_{2,m,1})=\Phi_{m-2}+k_m-1$, $\sum b(\Gamma_{3,m,1})=\Phi_{m-3}+t_{m-4}-k_{m-3}+1$. Moreover, $\Phi_m=\Phi_{m-1}+\Phi_{m-2}+\Phi_{m-3}+\frac{-3t_m+6t_{m-1}+t_{m-2}-1}{2}$ for $m\geq 7$. By induction, we can prove Property 4.3 below easily.

Property 4.3. *For $m\geq 4$, (1) $\Phi_m=\frac{m}{22}(-5t_m+14t_{m-1}+4t_{m-2})+\frac{1}{44}(67t_m-166t_{m-1}+5t_{m-2})+\frac{1}{4}$;*
(2) $\sum b(\Gamma_{1,m,1})=\frac{m}{22}(4t_m-9t_{m-1}+10t_{m-2})+\frac{1}{44}(19t_m+36t_{m-1}-169t_{m-2})-\frac{1}{4}$,
$\sum b(\Gamma_{2,m,1})=\frac{m}{22}(10t_m-6t_{m-1}-19t_{m-2})+\frac{1}{44}(-189t_m+156t_{m-1}+331t_{m-2})-\frac{1}{4}$,
$\sum b(\Gamma_{3,m,1})=\frac{m}{22}(-19t_m+29t_{m-1}+13t_{m-2})+\frac{1}{44}(237t_m-358t_{m-1}-157t_{m-2})+\frac{3}{4}$;
(3) $\sum_{j=4}^{m-1}\Phi_j=\frac{m}{44}(13t_m-10t_{m-1}+5t_{m-2})+\frac{2}{11}(-8t_m+8t_{m-1}-7t_{m-2})+\frac{m}{4}+2$;
(4) $B(\max\Gamma_{3,m,1})=\frac{m}{44}(-25t_m+48t_{m-1}+31t_{m-2})+\frac{1}{44}(173t_m-294t_{m-1}-213t_{m-2})+\frac{m+11}{4}$,
$B(\max\Gamma_{2,m,1})=\frac{m}{44}(-5t_m+36t_{m-1}-7t_{m-2})+\frac{1}{22}(-8t_m-69t_{m-1}+59t_{m-2})+\frac{m+10}{4}$,
$B(\max\Gamma_{1,m,1})=\frac{m}{44}(3t_m+18t_{m-1}+13t_{m-2})+\frac{1}{44}(3t_m-102t_{m-1}-51t_{m-2})+\frac{m+9}{4}$.

For any $n\geq 52$, let m such that $n\in[\Gamma_{3,m,1},\Gamma_{2,m,1},\Gamma_{1,m,1}]=[\frac{t_m+t_{m-2}-1}{2},\cdots,\frac{t_m+2t_{m-1}+t_{m-2}-3}{2}]$. We already determine the expression of $B(\max\Gamma_{j,m,1})$ for $j\in\{1,2,3\}$, $m\geq 5$. In order to calculate $B(n)$, we only need to calculate $\sum_{i=\min\Gamma_{j,m,1}}^{n}b(i)$.

Case 1. When $n\in\Gamma_{3,m,1}=[\frac{t_m+t_{m-2}-1}{2},\cdots,\frac{-t_m+4t_{m-1}+t_{m-2}-3}{2}]$ for $m\geq 7$. Denote

$$\begin{cases}\theta_m^1=\min\Gamma_{3,m,1}=\frac{t_m+t_{m-2}-1}{2};\\ \theta_m^2=\min\Gamma_{3,m,1}+|\Gamma_{3,m-3,1}|=\frac{-5t_m+10t_{m-1}+3t_{m-2}-1}{2};\\ \theta_m^3=\min\Gamma_{3,m,1}+|\Gamma_{3,m-3,1}|+|\Gamma_{2,m-3,1}|=\frac{-t_m+6t_{m-1}-3t_{m-2}-1}{2};\\ \eta_m^1=\min\Gamma_{3,m,1}+t_{m-4}-k_{m-3}+1=-2t_m+5t_{m-1}.\\ \theta_m^4=\max\Gamma_{3,m,1}+1=\min\Gamma_{2,m,1}=\frac{-t_m+4t_{m-1}+t_{m-2}-1}{2}.\end{cases}$$

Obviously, $\theta_m^3<\eta_m^1<\theta_m^4$ for $m\geq 7$, $\min\Gamma_{3,m,1}-\min\Gamma_{3,m-3,1}=t_{m-1}$. By Property 4.2, we have:

Property 4.4. *For $n\geq 52$, let m such that $n\in\Gamma_{3,m,1}$, then $m\geq 7$ and*

$$\sum_{i=\min\Gamma_{3,m,1}}^{n}b(i)$$
$$=\begin{cases}\sum_{i=\min\Gamma_{3,m-3,1}}^{n-t_{m-1}}b(i)+n-\min\Gamma_{3,m,1}+1, & \theta_m^1\leq n<\theta_m^2;\\ \sum_{i=\min\Gamma_{2,m-3,1}}^{n-t_{m-1}}b(i)+\sum b(\Gamma_{3,m-3,1})+n-\min\Gamma_{3,m,1}+1, & \theta_m^2\leq n<\theta_m^3;\\ \sum_{i=\min\Gamma_{1,m-3,1}}^{n-t_{m-1}}b(i)+\sum b(\Gamma_{3,m-3,1})+\sum b(\Gamma_{2,m-3,1})+n-\min\Gamma_{3,m,1}+1, & \theta_m^3\leq n<\eta_m^1;\\ \sum_{i=\min\Gamma_{1,m-3,1}}^{n-t_{m-1}}b(i)+\sum b(\Gamma_{3,m-3,1})+\sum b(\Gamma_{2,m-3,1})+\frac{-5t_m+10t_{m-1}-t_{m-2}+1}{2}, & otherwise.\end{cases}$$

Case 2. When $n\in\Gamma_{2,m,1}=[\frac{-t_m+4t_{m-1}+t_{m-2}-1}{2},\cdots,\frac{t_m+2t_{m-1}-t_{m-2}-3}{2}]$ for $m\geq 6$. Denote

$$\begin{cases}\theta_m^5=\min\Gamma_{2,m,1}+|\Gamma_{3,m-2,1}|=\frac{3t_m-5t_{m-2}-1}{2};\\ \eta_m^2=\min\Gamma_{2,m,1}+t_{m-3}-k_m+1=2t_{m-1}-t_{m-2}.\\ \theta_m^6=\min\Gamma_{3,m,1}+|\Gamma_{3,m-3,1}|+|\Gamma_{2,m-3,1}|=\frac{3t_m-2t_{m-1}-t_{m-2}-1}{2};\\ \theta_m^7=\max\Gamma_{2,m,1}+1=\min\Gamma_{1,m,1}=\frac{t_m+2t_{m-1}-t_{m-2}-1}{2}.\end{cases}$$

Obviously, $\theta_m^5<\eta_m^2\leq\theta_m^6$ for $m\geq 6$, $\min\Gamma_{2,m,1}-\min\Gamma_{3,m-2,1}=t_{m-1}$. By Property 4.2, we have:

Property 4.5. *For $n\geq 32$, let m such that $n\in\Gamma_{2,m,1}$, then $m\geq 6$ and*

$$\sum_{i=\min\Gamma_{2,m,1}}^{n}b(i)$$
$$=\begin{cases}\sum_{i=\min\Gamma_{3,m-2,1}}^{n-t_{m-1}}b(i), & \theta_m^4\leq n<\theta_m^5;\\ \sum_{i=\min\Gamma_{2,m-2,1}}^{n-t_{m-1}}b(i)+\sum b(\Gamma_{3,m-2,1}), & \theta_m^5\leq n<\eta_m^2;\\ \sum_{i=\min\Gamma_{2,m-2,1}}^{n-t_{m-1}}b(i)+\sum b(\Gamma_{3,m-2,1})+n-\eta_m^2+1, & \eta_m^2\leq n<\theta_m^6;\\ \sum_{i=\min\Gamma_{1,m-2,1}}^{n-t_{m-1}}b(i)+\sum b(\Gamma_{3,m-2,1})+\sum b(\Gamma_{2,m-2,1})+n-\eta_m^2+1, & otherwise.\end{cases}$$

Case 3. When $n \in \Gamma_{1,m,1} = [\frac{t_m+2t_{m-1}-t_{m-2}-1}{2}, \cdots, \frac{t_m+2t_{m-1}+t_{m-2}-3}{2}]$ for $m \geq 5$. Denote

$$\begin{cases} \theta_m^8 = \min\Gamma_{1,m,1} + |\Gamma_{3,m-1,1}| = \frac{t_m+3t_{m-2}-1}{2}; \\ \theta_m^9 = \min\Gamma_{1,m,1} + |\Gamma_{3,m-1,1}| + |\Gamma_{2,m-1,1}| = \frac{-t_m+4t_{m-1}+3t_{m-2}-1}{2}; \\ \eta_m^3 = \min\Gamma_{1,m,1} + t_{m-2} - k_m + 1 = 2t_{m-1}. \\ \theta_m^{10} = \max\Gamma_{1,m,1} + 1 = \min\Gamma_{3,m+1,1} = \frac{t_m+2t_{m-1}+t_{m-2}-1}{2}. \end{cases}$$

Obviously, $\theta_m^9 < \eta_m^3 < \theta_m^{10}$ for $m \geq 5$, $\min\Gamma_{1,m,1} - \min\Gamma_{3,m-1,1} = t_{m-1}$. By Property 4.2, we have:

Property 4.6. *For $n \geq 21$, let m such that $n \in \Gamma_{1,m,1}$, then $m \geq 5$ and*

$$\begin{aligned} &\sum_{i=\min\Gamma_{1,m,1}}^{n} b(i) \\ &= \begin{cases} \sum_{i=\min\Gamma_{3,m-1,1}}^{n-t_{m-1}} b(i), & \theta_m^7 \leq n < \theta_m^8; \\ \sum_{i=\min\Gamma_{2,m-1,1}}^{n-t_{m-1}} b(i) + \sum b(\Gamma_{3,m-1,1}), & \theta_m^8 \leq n < \theta_m^9; \\ \sum_{i=\min\Gamma_{1,m-1,1}}^{n-t_{m-1}} b(i) + \sum b(\Gamma_{3,m-1,1}) + \sum b(\Gamma_{2,m-1,1}), & \theta_m^9 \leq n < \eta_m^3; \\ \sum_{i=\min\Gamma_{1,m-1,1}}^{n-t_{m-1}} b(i) + \sum b(\Gamma_{3,m-1,1}) + \sum b(\Gamma_{2,m-1,1}) + n - \eta_m^3 + 1, & otherwise. \end{cases} \end{aligned}$$

Algorithm B. *Step 1. For $n \leq 51$, calculate $\sum_{i=1}^{n} b(i)$ by Property 4.2.*

Step 2. For $n \geq 52$, find the m and j such that $n \in \Gamma_{j,m,1}$, then $m \geq 7$. Calculate $B(\min\Gamma_{j,m,1}-1)$ by Property 4.3. And calculate $\sum_{i=\min\Gamma_{j,m,1}}^{n} b(i)$ by Property 4.4-4.6.

Step 3. $B(n) = B(\min\Gamma_{j,m,1} - 1) + \sum_{i=\min\Gamma_{j,m,1}}^{n} b(i)$.

Example. Consider $n = 60 \in \Gamma_{2,7,1} = [59, \cdots, 71]$. $\theta_7^4 = \frac{-t_7+4t_6+t_5-1}{2} = 59 \leq n < \theta_7^5 = \frac{3t_7-5t_5-1}{2} = 61$. By Property 4.5, $\sum_{i=\min\Gamma_{2,7,1}}^{60} b(i) = \sum_{i=\min\Gamma_{3,5,1}}^{60-t_6} b(i) = \sum_{i=15}^{16} b(i) = b(15) + b(16) = 2$. By Property 4.3, $B(\min\Gamma_{2,7,1} - 1) = B(\max\Gamma_{3,7,1}) = \frac{7}{44}(-25t_7 + 48t_6 + 31t_5) + \frac{1}{44}(173t_7 - 294t_6 - 213t_5) + \frac{18}{4} = 45$. Thus $B(60) = B(\max\Gamma_{3,7,1}) + \sum_{i=15}^{16} b(i) = 47$.

Now we turn to give the expressions of $B(t_m)$.

For $m \geq 7$, $\theta_m^8 \leq t_m < \theta_m^9$ and $\theta_{m-1}^6 \leq t_m - t_{m-1} < \theta_{m-1}^7$. By Property 4.5-4.6,

$$\begin{aligned} &\sum_{i=\min\Gamma_{1,m,1}}^{t_m} b(i) - \sum_{i=\min\Gamma_{1,m-3,1}}^{t_{m-3}} b(i) \\ =&\sum b(\Gamma_{3,m-1,1}) + \sum b(\Gamma_{3,m-3,1}) + \sum b(\Gamma_{2,m-3,1}) + 2t_m - 2t_{m-1} - 3t_{m-2} + 1 \\ =&\frac{m}{22}(-19t_m + 29t_{m-1} + 13t_{m-2}) + \frac{1}{44}(347t_m - 622t_{m-1} - 47t_{m-2}) + \frac{9}{4}. \end{aligned}$$

For $m \geq 4$, by induction, $\sum_{i=\min\Gamma_{1,m,1}}^{t_m} b(i)$ is equal to

$$\frac{m}{44}(23t_m - 38t_{m-1} - 3t_{m-2}) + \frac{1}{44}(-65t_m + 164t_{m-1} - 105t_{m-2}) + \frac{3m}{4} - \frac{9}{4}.$$

Since $\min\Gamma_{1,m,1} - 1 = \max\Gamma_{2,m,1}$, $B(t_m) = B(\max\Gamma_{2,m,1}) + \sum_{i=\min\Gamma_{1,m,1}}^{t_m} b(i)$. By Property 4.3, we can prove Theorem 21 in H.Mousavi and J.Shallit[6] in a novel way, i.e. **Theorem B**.

5 The number of distinct cubes, $C(n)$

Let ω be a factor with kernel K_m. By an analogous argument as Section 3, $\omega_p\omega_{p+1}\omega_{p+2} \prec \mathbb{T}$ has only one case: $G_p(K_m) = G_{p+1}(K_m) = G_1(K_m)$. Here $|G_1(K_m)| = t_m - k_m$, and

$$\omega\omega\omega = T_{m-1}[i, t_{m-1}]T_{m-2}T_{m-3}[1, t_{m-3} - 1]\underline{K_{m+4}}T_{m+1}[k_{m+4}, t_{m+1}]T_{m-3}T_{m-2}[1, i-1],$$

where $1 \leq i \leq k_{m+1} - 1$, $m \geq 3$, $Ker(\omega\omega\omega) = K_{m+4}$ and $|\omega| = t_m$.

Remark. By this case, we have that: all cubes in $\mathbb{T}$ are of length $3t_m$ for some $m \geq 3$. Furthermore, for all $m \geq 3$, there exists a cube of length $3t_m$ in $\mathbb{T}$. This is Theorem 7 in [6].

By the case of cubes, we define a set for $m \geq 7$,

$$\begin{aligned}\langle K_m, p\rangle =&\{P(\omega\omega\omega, p) : Ker(\omega\omega\omega) = K_m, |\omega| = t_{m-4}, \omega\omega\omega \prec \mathbb{T}\}\\ =&\{P(K_m, p) + \tfrac{-t_{m-2}+5t_{m-4}+1}{2}, \cdots, P(K_m, p) + t_{m-4} - 1\}.\end{aligned}$$

Then $\sharp\langle K_m, p\rangle = k_{m-3} - 1$. $P(K_m, 1) = \frac{t_m+t_{m-2}-1}{2}$, so $\langle K_m, 1\rangle = \{t_{m-1} + 2t_{m-4}, \cdots, \frac{3t_{m-1}-t_{m-3}-3}{2}\}$. Since $\langle K_m, 1\rangle$ are pairwise disjoint for different m, we get a chain $\langle K_7, 1\rangle, \langle K_8, 1\rangle, \cdots, \langle K_m, 1\rangle, \cdots$. Denote $c(n) = \sharp\{\omega\omega\omega : \omega\omega\omega \triangleright \mathbb{T}[1, n], \omega\omega\omega \not\prec \mathbb{T}[1, n-1]\}$, then $c(n) = 1$ if and only if $n \in \cup_{m\geq 7}\langle K_m, 1\rangle$.

Property 5.1. *$c(n) = 0$ for $n \leq 57$; for $n \geq 58$, let m such that $t_{m-1} + 2t_{m-4} \leq n < t_m + 2t_{m-3}$, then $m \geq 7$ and $c(n) = 1$ if and only if $n \leq t_{m-1} + k_{m+1} - 2$.*

Obviously, $C(n) = \sum_{i=1}^{n} c(i)$, and $C(n) = 0$ for $n \leq 57$. For $m \geq 7$, we denote

$$\hat{\alpha}_m = \min\langle K_m, 1\rangle = t_{m-1} + 2t_{m-4},\ \hat{\beta}_m = \max\langle K_m, 1\rangle = \tfrac{3t_{m-1}-t_{m-3}-3}{2}.$$

When $n \geq 58$, find m such that $\hat{\alpha}_m \leq n < \hat{\alpha}_{m+1}$. By Property 5.1 and the definition of Δ_m,

$$\begin{cases} C(\hat{\alpha}_m) = C(\min\langle K_m, 1\rangle) = \Delta_{m-4} + 1 = \frac{t_{m-6}+t_{m-7}-m+6}{2},\\ C(\hat{\beta}_m) = C(\max\langle K_m, 1\rangle) = \Delta_{m-3} = \frac{t_{m-5}+t_{m-6}-m+3}{2}.\end{cases}$$

Moreover, (1) when $\hat{\alpha}_m \leq n \leq \hat{\beta}_m$, $C(n) = C(\hat{\alpha}_m)+n-\hat{\alpha}_m$; (2) when $\hat{\beta}_m < n < \hat{\alpha}_{m+1}$, $C(n) = C(\hat{\beta}_m)$. Thus we get **Theorem C.**, i.e. the expression of $C(n)$ for all n.

For $m \geq 7$, since $t_m \geq \hat{\beta}_m = t_{m-1} + k_{m+1} - 2$, by Theorem C., we have $C(t_m) = C(\hat{\beta}_m)$, i.e.

Theorem 5.2. *For $m \leq 6$, $C(t_m) = 0$, for $m \geq 7$, $C(t_m) = \frac{t_{m-5}+t_{m-6}-m+3}{2}$.*

6 The number of repeated cubes, $D(n)$

For $m \geq 7$ and $p \geq 1$, we consider the vectors $\Gamma_{m,p} = [P(K_m, p), \cdots, P(K_m, p) + t_{m-1} - 1]$. Using Property 2.4, comparing minimal and maximal elements in these sets below, we have

$$\Gamma_{m,p} = [\Gamma_{m-3,P(a,p)+1}, \Gamma_{m-2,P(b,p)+1}, \Gamma_{m-1,P(c,p)+1}] \text{ for } m \geq 10.$$

Thus we establish the recursive relations for all $\Gamma_{m,p}$, $m \geq 10$. We arrange all elements of $\langle K_m, p\rangle$ in ascending order as vector $[\langle K_m, p\rangle]$. It is easy to check that

$$\Gamma_{m,p} = [\underbrace{P, \cdots, P + \tfrac{-t_{m-2}+5t_{m-4}-1}{2}}_{\frac{-t_{m-2}+5t_{m-4}+1}{2}}, \underbrace{[\langle K_m, p\rangle]}_{\frac{t_{m-2}-3t_{m-4}-1}{2}}, \underbrace{P + t_{m-4}, \cdots, P + t_{m-1} - 1}_{t_{m-2}+t_{m-3}}].$$

Here we denote $P(K_m, p)$ by P for short. The number under "$\smile$" means the number of elements.

By an analogous argument in Section 4, we get the recursive structure of the positions of repeated cubes in $\mathbb{T}$, called cube trees. Here π is a substitution over $\{\langle K_m, p\rangle : m \geq 7, p \geq 1\}$ that

$$\begin{cases}\pi\langle K_8, p\rangle = \langle K_7, P(a, p) + 1\rangle;\ \pi\langle K_9, p\rangle = \langle K_7, P(b, p) + 1\rangle \cup \langle K_8, P(a, p) + 1\rangle;\\ \pi\langle K_m, p\rangle = \langle K_{m-3}, P(a, p) + 1\rangle \cup \langle K_{m-2}, P(b, p) + 1\rangle \cup \langle K_{m-1}, P(c, p) + 1\rangle, m \geq 10.\end{cases}$$

The cube trees contain all $\langle K_m, p\rangle$, i.e. the positions of all cubes in $\mathbb{T}$.

Let $p = 1$, $\Gamma_{m,1} = [P(K_m, 1), \cdots, P(K_{m+1}, 1) - 1] = [\frac{t_m+t_{m-2}-1}{2}, \cdots, \frac{t_{m+1}+t_{m-1}-3}{2}]$.

Denote $d(n) = \sharp\{(\omega, p) : \omega_p\omega_{p+1}\omega_{p+2} \triangleright \mathbb{T}[1, n]\}$ the number of cubes ending at position n. By Property 6.1 below, we can calculate $d(n)$, and obversely calculate $D(n)$ by $D(n) = \sum_{i=1}^{n} d(i)$.

Property 6.1. *For $n \le 51$, $d(n) = 0$; and*

$$\begin{cases} d(\Gamma_{7,1}) = d([52,\cdots,95]) = [\underbrace{0,\cdots,0}_{6},1,\underbrace{0,\cdots,0}_{37}]; \\ d(\Gamma_{8,1}) = d([96,\cdots,176]) = [\underbrace{0,\cdots,0}_{11},1,1,\underbrace{0,\cdots,0}_{30},1,\underbrace{0,\cdots,0}_{37}]; \\ d(\Gamma_{9,1}) = d([177,\cdots,325]) = [\underbrace{0,\cdots,0}_{20},\underbrace{1,\cdots,1}_{4},\underbrace{0,\cdots,0}_{6},1,\underbrace{0,\cdots,0}_{48},1,1,\underbrace{0,\cdots,0}_{30},1,\underbrace{0,\cdots,0}_{37}]; \\ d(\Gamma_{m,1}) = [d(\Gamma_{m-3,1}), d(\Gamma_{m-2,1}), d(\Gamma_{m-1,1})] + [\ \underbrace{0,\cdots,0,}_{\frac{-t_{m-2}+5t_{m-4}+1}{2}}\ \underbrace{1,\cdots,1,}_{\frac{t_{m-2}-3t_{m-4}-1}{2}}\ \underbrace{0,\cdots,0}_{t_{m-2}+t_{m-3}}\ ], m \ge 10. \end{cases}$$

Now we turn to give a fast algorithm. Denote the sum of all elements in vector $d(\Gamma_{m,1})$ by $\sum d(\Gamma_{m,1})$. An immediately corollaries of Property 6.1 are $\sum d(\Gamma_{m,1}) = \sum d(\Gamma_{m-3,1}) + \sum d(\Gamma_{m-2,1}) + \sum d(\Gamma_{m-1,1}) + \frac{t_{m-2}-3t_{m-4}-1}{2}$. By induction, we can prove Property 6.2 below easily.

Property 6.2. *For $m \ge 7$, (1) $D(\min \Gamma_{m+1,1}) = D(\max \Gamma_{m,1})$;*
(2) $\sum d(\Gamma_{m,1}) = \frac{m}{22}(7t_m - 13t_{m-1} + t_{m-2}) + \frac{1}{44}(-41t_m + 74t_{m-1} - 7t_{m-2}) + \frac{1}{4}$;
(3) $D(\max \Gamma_{m,1}) = \sum_{j=7}^{m} \sum d(\Gamma_{j,1}) = \frac{m}{44}(9t_m - 12t_{m-1} - 5t_{m-2}) + \frac{3}{11}(-2t_m + 2t_{m-1} + t_{m-2}) + \frac{m}{4}$.

For any $n \ge 52$, let m such that $n \in \Gamma_{m,1}$. Now we calculate $\sum_{i=\min \Gamma_{m,1}}^{n} d(i)$. Denote

$$\begin{cases} \hat{\theta}_m^1 = \min \Gamma_{m,1} = \frac{t_m + t_{m-2} - 1}{2}; \\ \hat{\eta}_m^1 = \min \Gamma_{m,1} + \frac{-t_{m-2}+5t_{m-4}+1}{2} = \frac{t_m + 5t_{m-4}}{2}; \\ \hat{\eta}_m^2 = \min \Gamma_{m,1} + \frac{-t_{m-2}+5t_{m-4}+1}{2} + \frac{t_{m-2}-3t_{m-4}-1}{2} = \frac{3t_{m-1}-t_{m-3}-1}{2}; \\ \hat{\theta}_m^2 = \min \Gamma_{m,1} + |\Gamma_{m-3,1}| = \frac{3t_{m-1}-t_{m-3}-1}{2} = \hat{\eta}_m^2; \\ \hat{\theta}_m^3 = \min \Gamma_{m,1} + |\Gamma_{m-3,1}| + |\Gamma_{m-2,1}| = \frac{3t_{m-1}+t_{m-3}-1}{2}. \end{cases}$$

Obviously, $\hat{\theta}_m^1 < \hat{\eta}_m^1 < \hat{\eta}_m^2 = \hat{\theta}_m^2 < \hat{\theta}_m^3 < \hat{\theta}_{m+1}^1$ for $m \ge 7$. By Property 6.1, we have:

Property 6.3. *For $n \ge 326$, let m such that $n \in \Gamma_{m,1}$, then $m \ge 10$ and*

$$\begin{aligned} &\sum_{i=\min \Gamma_{m,1}}^{n} d(i) \\ &= \begin{cases} \sum_{i=\min \Gamma_{m-3,1}}^{n-t_{m-1}} d(i), & \hat{\theta}_m^1 \le n < \hat{\eta}_m^1; \\ \sum_{i=\min \Gamma_{m-3,1}}^{n-t_{m-1}} d(i) + n - \frac{t_m + 5t_{m-4}}{2} + 1, & \hat{\eta}_m^1 \le n < \hat{\theta}_m^2; \\ \sum_{i=\min \Gamma_{m-2,1}}^{n-t_{m-1}} d(i) + \sum d(\Gamma_{m-3,1}) + \frac{t_{m-2}-3t_{m-4}-1}{2}, & \hat{\theta}_m^2 \le n < \hat{\theta}_m^3; \\ \sum_{i=\min \Gamma_{m-1,1}}^{n-t_{m-1}} d(i) + \sum d(\Gamma_{m-3,1}) + \sum d(\Gamma_{m-2,1}) + \frac{t_{m-2}-3t_{m-4}-1}{2}, & otherwise. \end{cases} \end{aligned}$$

Algorithm D. *Step 1. For $n \le 325$, calculate $\sum_{i=1}^{n} d(i)$ by Property 6.1.*

Step 2. For $n \ge 326$, find the m such that $n \in \Gamma_{m,1}$, then $m \ge 10$. Calculate $\sum_{i=\min \Gamma_{m,1}}^{n} d(i)$ by Property 6.3. And calculate $D(\min \Gamma_{m,1} - 1) = D(\max \Gamma_{m-1,1})$ by Property 6.2.

Step 3. $D(n) = D(\max \Gamma_{m-1,1}) + \sum_{i=\min \Gamma_{m,1}}^{n} d(i)$.

Example. Consider $n = 500 \in \Gamma_{10,1} = [326,\cdots,599]$. Since $\hat{\theta}_{10}^3 = \frac{3t_9+t_7-1}{2} = 451 \le n < \hat{\theta}_{11}^1 = \frac{t_{11}+t_9-1}{2} = 600$, by Property 6.3, $\sum_{i=\min \Gamma_{10,1}}^{500} d(i) = \sum_{i=\min \Gamma_{9,1}}^{500-t_9} d(i) + \sum d(\Gamma_{7,1}) + \sum d(\Gamma_{8,1}) + \frac{t_8-3t_6-1}{2}$. Here $\sum_{i=\min \Gamma_{9,1}}^{500-t_9} d(i) = \sum_{i=177}^{226} d(i) = d([197,198,199,200]) + d(207) = 5$; $\sum d(\Gamma_{7,1}) = 1$, $\sum d(\Gamma_{8,1}) = 3$ and $\frac{t_8-3t_6-1}{2} = 8$. Thus $\sum_{i=\min \Gamma_{10,1}}^{500} d(i) = 17$. By Property 6.2, $D(\max \Gamma_{9,1}) = \frac{9}{44}(9t_9 - 12t_8 - 5t_7) + \frac{3}{11}(-2t_9 + 2t_8 + t_7) + \frac{9}{4} = 12$. Thus $D(500) = D(\max \Gamma_{9,1}) + \sum_{i=\min \Gamma_{10,1}}^{500} d(i) = 12 + 17 = 29$.

In fact, the positions of the last letters of the 29 cubes are $\{58, 107, 108, 139, 197, 198, 199, 200, 207, 256, 257, 288, 332, 362, 363, 364, 365, 366, 367, 368, 369, 381, 382, 413, 471, 472, 473, 474, 481\}$.

Now we turn to give the expressions of $D(t_m)$.
For $m \geq 10$, $\hat{\theta}^3_m \leq t_m < \hat{\theta}^1_{m+1}$ and $\hat{\theta}^2_{m-1} \leq t_m - t_{m-1} < \hat{\theta}^3_{m-1}$. By Property 6.3,

$$\begin{aligned}
&\textstyle\sum_{i=\min\Gamma_{m,1}}^{t_m} d(i) - \sum_{i=\min\Gamma_{m-3,1}}^{t_{m-3}} d(i) \\
=&\textstyle\sum d(\Gamma_{m-3,1}) + \sum d(\Gamma_{m-2,1}) + \frac{t_{m-2}-3t_{m-4}-1}{2} + \sum d(\Gamma_{m-4,1}) + \frac{t_{m-3}-3t_{m-5}-1}{2} \\
=&\textstyle\sum d(\Gamma_{m-1,1}) + \frac{t_{m-2}-3t_{m-4}-1}{2} \\
=&\textstyle\frac{m}{22}(7t_{m-1} - 13t_{m-2} + t_{m-3}) + \frac{1}{44}(-121t_{m-1} + 188t_{m-2} + 57t_{m-3}) - \frac{1}{4} \\
=&\textstyle\frac{m}{22}(t_m + 6t_{m-1} - 14t_{m-2}) + \frac{1}{44}(57t_m - 178t_{m-1} + 131t_{m-2}) - \frac{1}{4}.
\end{aligned}$$

For $m \geq 7$, by induction, $\sum_{i=\min\Gamma_{m,1}}^{t_m} d(i)$ is equal to

$$\begin{aligned}
\textstyle\sum_{i=\min\Gamma_{m,1}}^{t_m} d(i) =& \textstyle\frac{m}{44}(-7t_m + 2t_{m-1} + 21t_{m-2}) + \frac{1}{11}(-10t_m + 21t_{m-1} - 6t_{m-2}) - \frac{m}{12} \\
&+ 0[m \equiv 0 (\mathrm{mod}\ 3)] + \textstyle\frac{1}{3}[m \equiv 1 (\mathrm{mod}\ 3)] + \frac{2}{3}[m \equiv 2 (\mathrm{mod}\ 3)].
\end{aligned}$$

Here $[P]$ is Iverson notation, and equals 1 if P holds and 0 otherwise.

By Property 6.2 and $D(t_m) = D(\max\Gamma_{m-1,1}) + \sum_{i=\min\Gamma_{m,1}}^{t_m} d(i)$, we prove Theorem 21 in H.Mousavi and J.Shallit[6] in a novel way, i.e. **Theorem D**.

References

[1] J. Justin, G. Pirillo, On a characteristic property of Arnoux-Rauzy sequences. *RAIRO: Theoret. Informatics Appl.* **36.4** (2002) 385-388.

[2] A. Glen, *On Sturmian and Episturmian Words, and Related Topics,* Ph.D. thesis. The University of Adelaide, Australia (2006).

[3] Y.K. Huang, Z.Y. Wen, Kernel words and gap sequence of the Tribonacci sequence. *Acta Mathematica Scientia (Series B).* **36.1** (2016) 173-194.

[4] Y.K. Huang, Z.Y. Wen, The number of distinct and repeated squares and cubes in the Fibonacci sequence. arXiv:1603.04211.

[5] Y.K. Huang, Z.Y. Wen, The numbers of repeated palindromes in the Fibonacci and Tribonacci sequences. arXiv:1604.05021.

[6] H. Mousavi, J. Shallit, Mechanical proofs of properties of the Tribonacci word, in *Combinatorics on Words*. Springer International Publishing. (2014) 170-190.

[7] B. Tan, Z.Y. Wen, Some properties of the Tribonacci sequence. *European J Combin.* **28** (2007) 1703-1719.